

(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 321 811 B2

(12) **NEW EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the opposition decision:
22.12.1999 Bulletin 1999/51

(51) Int. Cl.⁶: **A21D 8/04**

(45) Mention of the grant of the patent:
17.06.1992 Bulletin 1992/25

(21) Application number: 88120669.2

(22) Date of filing: 10.12.1988

(54) **Method for improving flour dough**
Methode zur Verbesserung von Mehlteig
Méthode pour l'amélioration de la pâte de farine

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(30) Priority: 21.12.1987 US 136003

(43) Date of publication of application:
28.06.1989 Bulletin 1989/26

(73) Proprietor: DSM N.V.
6411 TE Heerlen (NL)

(72) Inventors:
• Haarasilta, Sampsa
SF-04230 Kerava (FI)
• Väisänen, Seppo
SF-04260 Kerava (FI)
• Scott, Don
Schaumburg, Ill. (US)

(74) Representative:
Matulewicz, Emil Rudolf Antonius et al
DSM- Patents & Trademarks (994-0760),
P.O. Box 1
2600 MA Delft (NL)

(56) References cited:
US-A- 2 783 150 US-A- 4 632 905
US-A- 4 894 340

- BROT UND GEBÄCK, vol. 9, no. 6, June 1955, pages 101-103; P. MALTHA: "Weitere Untersuchungen über die Backverbesserung durch 1-Ascorbinsäure"
- CHEMICAL ABSTRACTS, vol. 71, no. 11, 24th November 1969, pages 225,226, no. 100605k, Columbus, Ohio, US; M.V. POLYAK et al.: "Glucose oxidase as a conditioner in bread baking", & FERMENTY MED., PISHCH. PROM. SEL. KHOZ. 1968, 155-7
- CEREAL CHEMISTRY, vol. 64, no. 3, May-June 1987, pages 172-176, St. Paul, MN, US; S.P. KAUFMAN et al.: "Evaluation of sulfhydryl oxidase as a strengthening agent for wheat flour dough"
- Young and Nimmo, Proceedings of the Biochemical Society, 1972
- Motte et al, Biochemistry, vol. 26, November 1987, pp. 7363-7371
- Don Scott, Enzymes in Food Processing, 1975, pp. 222-229
- Kaufmann and Fennema, Cereal Chemistry, 64(3), 1987, pp. 172-176
- Novo's Handbook of Practical Biotechnology, 1986, pp. 96-99
- Fluka catalogue
- Statement regarding deposit date of *Aspergillus niger* ATCC 1004 and Strain Descriptions
- Ullmann's Encyclopedia of Industrial Chemistry, Fifth, completely revised Edition, Vol. A4, Germany, 1985, pp. 331-347

EP 0 321 811 B2

Description

[0001] This invention relates to a method for improving the rheological properties of a flour dough which comprises combining flour, yeast, water and an effective amount of an enzyme preparation comprising sulfhydryl oxidase and glucose oxidase and mixing said ingredients to form a suitable baking dough. The invention results in stronger doughs with improved rheological properties as well as a final baked product with improved texture.

[0002] The "strength" or "weakness" of dough is an important aspect of baking. Flours with a low protein content are customarily characterized as "weak"; the gluten (the cohesive, extensible, rubbery mass which is formed by mixing flour and water) formed with weak flour will be very extensible under stress, but will not return to its original dimensions when the stress is removed. Flours with a high protein content are customarily characterized as "strong" and the gluten formed with strong flour will be less extensible than a weak flour, and stress which is applied during mixing will be restored without breakdown to a greater extent than a weak flour. Strong dough is generally preferred in most baking contexts because of the superior rheological and handling properties of the dough and the superior form and texture qualities of the final baked product made from the dough.

[0003] For example, stronger dough is generally more stable; the stability of dough is one of the most important (if not the most important) characteristics of baking dough.


[0004] American Association of Cereal Chemists Method 36-01A defines dough stability as "(a) the range of dough time over which a positive Response is obtained; and (b) that property of a rounded dough by which it resists flattening under its own weight over a course of time". Response is defined, by the same Method, as "the reaction of dough to a known and specific stimulus, substance or set of conditions, usually determined by baking it in comparison with a control".

[0005] Stable dough is particularly useful in large scale applications where it may be difficult to control all processing parameters; strong dough will exhibit a greater tolerance of, e.g. mixing time and proofing time, and still result in quality products. Less stable dough will exhibit less tolerance in this regard.

[0006] Bakers have long used dough "conditioners" to strengthen the dough. It is suggested that such conditioners, which consist primarily of non-specific oxidants such as bromates, peroxides, iodates and ascorbic acid, help form inter-protein bonds which strengthen the dough. However, non-specific oxidants have numerous drawbacks; in particular, they can have a negative effect on the organoleptic qualities of the final product and are relatively expensive in commercial quantities and, in the case of bromates, are not permitted in certain countries.

[0007] The use of enzymes as dough conditioners has been considered as an alternative to non-specific oxidants. In particular, glucose oxidase has been used - sometimes in combinations with other conditioners - to condition or "mature" flour. U.S. Patent No. 2,783,150 (Luther) discusses the treatment of flour with glucose oxidase with allegedly can be used to form an improved dough with better handling properties and a high quality final baked product. However, the effects of glucose oxidase are somewhat contradictory. Water absorption of the dough is increased but glucose oxidase, in some contexts, may actually impair dough rheology and has never been successfully used as a dough conditioner.

[0008] It has also been suggested that the enzyme sulfhydryl oxidase could be used to strengthen dough. Sulfhydryl oxidase ("SHX") catalyzes - in the presence of oxygen - the conversion of thiol compounds to their corresponding disulfides according to the equation:



[0009] The role played by sulfur containing reactive groups in wheat proteins has not been fully defined but it is suggested that the reaction of free sulfhydryl groups to form disulfide bonds has an important role in the mixing and strength of dough. In particular, if disulfide bonds are formed between two protein chains, the resulting cross-linking of chains could strengthen the dough. Hence, SHX could be expected to strengthen dough by catalyzing the reaction of free sulfhydryl groups into inter-protein disulfide bonds.

[0010] However, Kaufmann et al., Cereal Chemistry 64:3 (1987), evaluated bovine SHX's ability to strengthen wheat dough and concluded that it did not have any strengthening effect. The baking tests reported by Kaufmann et al. did not show any "noticeable" effect of SHX on loaf volume, and mixograph studies on SHX treated dough which did not show any "noticeable" effect on the time to reach a mixing peak or the extent of dough breakdown. Kaufmann et al. also evaluated the effect of SHX on flour/buffer suspensions and concluded that SHX did not show any effect on the free-SH groups of flour. Kaufmann et al. state that - for a number of possible reasons - SHX was not able to catalyze formation of disulfide bonds in the systems tested.

[0011] It has now been discovered, however, that inclusion of an enzyme preparation which comprises glucose oxidase and microbial SHX into a flour, water and yeast mix appreciably and significantly strengthens the resulting dough; the dough exhibits improved rheological qualities, and in particular, demonstrates increased stability. The final, baked product made from such dough also exhibits improved form and texture qualities.

EP 0 321 811 B2

assess general quality of flour and its responses to improving agents*.

[0025] The farinograph method determines the water intake of a particular flour and the mixing tolerance of the resulting dough. Better baking flours, and dough, will exhibit higher farinograph values. If a particular flour shows relatively high water intake, and the mixing tolerance of the resulting dough is good, the farinograph curve shows retention of most if not all of the initial height over time. The machinability and baking quality of such a dough is likely to be excellent.

[0026] AACC Method 54-21 defines the farinograph as follows: "the farinograph measures and records resistance of a dough to mixing. It is used to evaluate absorption of flours and to determining stability and other characteristics of doughs during mixing".

[0027] Baking conditions used for baking bread from the dough prepared as above were as follows: oven: normal hearth oven (Dahlien) w/10 seconds steaming

flour time: 30 minutes
 final proofing: 30 minutes/37°C./75 % humidity
 baking time: 25 minutes at 220°C.
 cooling time: 1 hour/20°C.

TABLE I

Comparison of Rheological Properties of Doughs					
ENZYME ADDITION	Dough Sample				
	1	2	3	4	5
SHX Units/kg flour	-	38	77	154	770
GO Units/kg flour	-	110	219	438	2190
CAT Units/kg flour	-	2	4	7	40
SHX/GO Unit ratio	-	0.34	0.35	0.35	0.35
GO/CAT Unit ratio	-	55	55	55	55
RESULTS					
Farinogram Water absorption %	62	62	62	62	62
Development of the dough/(measured in minutes)	2	3	3.5	2	2
Stability/(measured in minutes)	3	12	23	19	2
Softening/(after?) 12 min Brabender Units (B.u.)	50	30	10	10	120
Resistance to Extens. Brabender Units (B.u.)	420	450	580	610	980
Extension,mm Brabender Units (B.u.)	180	165	143	139	86
Ratio resistance/extens B.u./mm	2.4	2.7	4.1	4.4	11
Maximum extension (B.u.)	620	640	760	720	980

[0028] Table I, above sets forth rheological properties for four doughs, a control dough (dough #1) and doughs conditioned with the enzyme preparations having different levels of SHX, glucose oxidase and catalase (doughs #2, #3, #4 and #5). The data demonstrates that dough mixed with an enzyme preparation containing glucose oxidase and SHX exhibit significantly improved rheological properties when compared with the control dough. In particular, dough#3 (23 minutes) and dough#4 (19 minutes) exhibit dramatic increases in stability compared to the control dough (3 minutes). The improved stability of the doughs treated with the enzyme preparation indicates that such doughs will exhibit better handling and machinability properties. The analysis data for the doughs treated with the enzyme preparation also suggests that these doughs were significantly strengthened. The resistance to extension, the maximum extension and the ratio of resistance to extension/extension all indicate that the treated doughs were significantly strengthened.

[0029] As noted, doughs with improved stability and strength generally also result in final baked products with improved qualities. Baking tests confirmed that doughs treated with the method of the present invention provided superior final products.

TABLE II

Results of Baking Tests			
	Loaf Sample		
	1	2	3
Dough sample (Table I)	1 (control)	3	4
Dough consistency			
After mixing (B.u.)	325	315	320
After floor time (B.u.)	285	290	300
Loaf weight (g)	370	370	370
Loaf height (mm)	76	79	76
Loaf width (mm)	172	171	173
Loaf H/W ratio	0.44	0.46	0.44
Loaf volume (ml)	1230	1340	1290
Loaf Spec. Vol. (ml/kg)	3310	3620	3490
Loaf moisture (%)	45.2	45.0	45.0

[0030] Table II sets forth baking results for loafs baked from doughs # 1 (control), # 3 and # 4 referred to in Table I. The control dough (loaf # 1) was not treated with any enzyme preparations and dough samples # 3 and # 4 (loaf samples # 2 and # 3 respectively) were treated with the enzyme preparation as set forth in Table I. The data from the baking tests set forth in Table II demonstrates that - compared to the control dough - doughs treated with the SHX/glucose oxidase enzyme preparation exhibited improved size and texture. In particular, loaf sample # 2 (baked with dough sample # 3) exhibited higher loaf volume (1340 ml. versus 1230 ml), higher specific volume (3620 ml/kg versus 3310 ml/kg) than the control sample. These height and width values demonstrate that loaf samples # 2 and # 3 were rounder and more symmetrical in shape, evidence of greater dough strength. In addition, the porosity of these loafs were more uniform meaning that the pores are the same size both near the crust and the center of the loafs.

[0031] Organoleptic comparison of the three loaf samples indicated that loaf samples # 2 and # 3 demonstrated improved texture properties compared to the control.

[0032] The baking results suggest that the present invention will help bakers achieve a larger loaf volume. In the commercial context this means that bakers could use wheat with a lower protein content, which is cheaper, to achieve the desired loaf size and/or can utilize a smaller dough plug to achieve the desired loaf size; both possibilities could potentially result in substantial savings in material costs to the baker.

[0033] In order to determine the effect of varying levels of SHX and glucose oxidase in enzyme preparations used to treat baking doughs, the following enzyme samples were developed:

Enzyme Sample A (prepared from A. niger cells) Activities:

GO 1.0 U/mg
SHX 11.5 U/mg
CAT -

Enzyme Sample B (prepared from A. niger cells) Activities:

GO 129 U/mg
SHX 0.4 U/mg
CAT 0.2 U/mg

Enzyme Sample C (prepared from A. niger cells) Activities:

GO 122 U/mg

SHX 0.8 U/mg
 CAT 2.0 U/mg

[0034] Tables III and IV below set forth data regarding the rheological properties of doughs prepared with Enzyme Samples A, B and C as well as a control dough.

TABLE III: Rheological Properties of Dough Samples - Various Enzyme Levels

Enzyme Sample	Activities		Water Absorp %	Dough Devel min	Dough Stabli min	Dough Soft B.u.	Resist B.u.	Ext min	Res/Ext ratio	Max B.u.
	GO	SHX								
Control			62	1.5	2.0	50	420	180	2.4	620
Enzyme Sample A	4	38	62	1.5	2.5	50	470	176	2.7	640
	8	77	62	1.5	2.0	60	470	174	2.7	640
Enzyme Sample B	645	2	62	2	3.5	20	640	125	5.1	740
	1290	4	62	2	15	10	640	126	5.1	760
	2580	8	62	2	14.5	20	660	125	5.3	760
	12900	40	62	2	2.5	120	980	95	10	980
Enzyme Sample C			62	2	3	30	720	129	5.6	900

[0035] The doughs set forth in Table III were prepared with "weak" Finnish wheat flour that had been treated with ascorbic acid (a non-specific oxidant) and preparation as described above. Although this data suggests, as indicated by the prior art, that glucose oxidase alone can have a conditioning effect, relatively large (and uneconomical) quantities of glucose oxidase are required to achieve appreciable strengthening. The data also suggests that a glucose oxidase/SHX combination is the most efficient and economical preparation for dough conditioning.

TABLE IV: Rheological Properties of Dough Samples

Enzyme Preparations	GO		Activities		Water Absorp %	Dough Devel min	Dough Stabill min	Dough Soft B.u.	Resist. B.u.	Ext. min	Res/Ext ratio	Max B.u.
	GO	SHX	Catalase	Catalase								
Amer. Bromated Flour (strong)	-	-	-	-	60	2.5	10	40	330	261	1.3	620
	110	38	2	2	60	2	15	0	400	210	1.9	720
	219	77	4	4	60	2	15	0	670	175	3.8	900
	440	154	7	7	60	5	13	0	735	168	4.4	860
Amer. Unbromated Flour (strong)	-	-	-	-	61	2.5	8	40	340	224	1.5	640
	110	38	2	2	61	2.5	12	20	390	188	2.1	580
	219	77	4	4	61	2	15	0	525	178	3.0	720
	440	154	7	7	61	5	12	0	610	182	3.4	720
Amer. (Weak) +SHX (x2620)	-	-	-	-	57	1.5	4	90	280	172	1.6	390
	110	38	2	2	57	1.5	3	100	370	149	2.5	440
	219	77	4	4	57	1.5	3.5	70	430	137	3.1	480
	440	154	7	7	57	1.5	4	60	515	140	3.7	560

[0036] Table IV sets forth data for doughs prepared using different US flours, both strong and weak. Compared to the control samples, doughs prepared with these flours that were treated with an SHX/glucose oxidase enzyme preparation generally demonstrated increased stability and strength, as indicated by the resistance and extension data. The effect was greater in the case of the "strong" flours (bromated and unbromated) than for the unbromated "weak" flours.

[0037] Table V set below sets forth data regarding rheological properties of doughs prepared with Finnish rye meal. A control dough with no enzyme treatment (sample # 1) and dough samples with varying levels of SHX and glucose oxidase were prepared.

TABLE V

Rheological Properties of Rye Dough* Samples								
Enzyme Preparations	Activities			Water Absorp %	Dough Devel min	Dough Stabil min	Dough Soft B.u.	
	GO	SHX	Catalase					
O - Control				75	3.5	3	20	Au°C
1.	110	38	2	73	3	>12	0	
2.	219	77	4	73	3	>12	0	
3.	440	154	7	74	3	>12	+10	
4.	645	2	1	74	3.5	>12	0	
5.	1290	4	2	74	3.5	>12	+10	
6.	2580	8	4	75	3.5	>12	+10	
7.	12900	40	20	74	3.5	6	20	

*Analysis of the rye meal samples:

Ash content 1.74 %
 Falling number 125
 Amylogram 260 B.U. at 66°C

[0038] The data in Table V shows that an enzyme preparation with SHX/GO acts as a dough conditioner; the most efficient - and cost effective - preparations (samples 1-3) will probably be more effective in large scale contexts.

[0039] The results of these experiments demonstrate that use of an enzyme preparation containing SHX and glucose oxidase appreciably and significantly improves the rheological properties of dough. The effect of enzyme preparation is more pronounced with certain flours and may be enhanced by the presence of a dough conditioner such as ascorbic acid. It is believed that oxidative enzymes and non-specific oxidants act synergistically.

[0040] The foregoing general discussion and experimental examples are intended to be illustrative of the present invention, and are not to be considered as limiting. Other variations within the spirit and scope of this invention are possible and will present themselves to those skilled in the art.

45 Claims

1. A method for improving rheological properties of a flour dough characterised in that flour, yeast and water are combined with an effective amount of a microbial enzyme preparation comprising sulphhydryl oxidase and glucose oxidase and said ingredients are mixed to form a suitable baking dough, the ratio of sulphhydryl oxidase to glucose oxidase in said preparation, based on units of enzymes present, being in the range of 0.003 to 10 and sulphhydryl oxidase being present in said dough in an amount of 35-800 units per kg of flour.
2. A method in accordance with claim 1 characterised in that the enzyme preparation is derived from Aspergillus niger cells.
3. A method in accordance with claim 1 characterised in that the enzyme preparation is added in an amount of about 80 units of sulphhydryl oxidase per kilogram of flour.

EP 0 321 811 B2

4. A method in accordance with claim 1 characterised in that the flour is wheat flour.
5. A method in accordance with claim 1 characterised in that the sulphhydryl oxidase and glucose oxidase are present in said preparation in a ratio of about 0.35, based on units of sulphhydryl oxidase and glucose oxidase present.
6. A method in accordance with claim 1 characterised in that the flour is fortified by non-specific dough conditioners.
7. A method in accordance with claim 6 characterised in that the flour is fortified with ascorbic acid as a dough conditioner.

Patentansprüche

1. Methode zur Verbesserung der rheologischen Eigenschaften eines Mehlteiges, dadurch gekennzeichnet, dass Mehl, Hefe und Wasser mit einer wirksamen Menge eines microbiellen Enzympräparates zusammengegeben werden, welches Sulphydryl-Oxidase und Glucose-Oxidase enthält, und dass diese Zutaten gemischt werden, um einen geeigneten Backteig zu bilden, wobei das Verhältnis von Sulphydryl-Oxidase zu Glucose-Oxidase im Präparat, basierend auf den anwesenden Enzym-Einheiten, im Bereich von 0.003 bis 10 liegt, und die Sulphydryl-Oxidase im Teig in einer Menge von 35-800 Einheiten pro kg Mehl vorhanden ist.
2. Methode gemäss Anspruch 1, dadurch gekennzeichnet, dass das Enzympräparat aus Aspergillus niger Zellen gewonnen wird.
3. Methode gemäss Anspruch 1, dadurch gekennzeichnet, dass das Enzympräparat in einer Menge von etwa 80 Einheiten an Sulphydryl-Oxidase pro Kilogramm Mehl zugegeben wird.
4. Methode gemäss Anspruch 1, dadurch gekennzeichnet, dass das Mehl Weizenmehl ist.
5. Methode gemäss Anspruch 1, dadurch gekennzeichnet, dass die Sulphydryl-Oxidase und Glucose-Oxidase im genannten Präparat in einem Verhältnis von etwa 0.35, basierend auf den Einheiten an Sulphydryl-Oxidase und Glucose-Oxidase, vorhanden sind.
6. Methode gemäss Anspruch 1, dadurch gekennzeichnet, dass das Mehl durch nicht-spezifische TeigKonditioniermittel gestärkt ist.
7. Methode gemäss Anspruch 6, dadurch gekennzeichnet, dass das Mehl mit Ascorbinsäure als Konditioniermittel gestärkt ist.

Revendications

1. Procédé pour améliorer les propriétés rhéologiques d'une pâte de farine, caractérisé en ce que de la farine, de la levure et de l'eau sont combinées avec une quantité efficace d'une préparation d'enzyme microbienne comprenant de la sulphydryle oxydase et de la glucose oxydase, et en ce que lesdits ingrédients sont mélangés pour former une pâte à cuire convenable, le rapport entre la sulphydryle oxydase et la glucose oxydase, sur la base des unités d'enzyme présentes, étant compris dans la gamme allant de 0,003 à 10, et la sulphydryle oxydase étant présente dans ladite pâte dans une quantité allant de 35 à 800 unités par kg de farine.
2. Procédé selon la revendication 1, caractérisé en ce que la préparation d'enzyme est dérivée de cellules d'Aspergillus niger.
3. Procédé selon la revendication 1, caractérisé en ce que la préparation d'enzyme est ajoutée dans une quantité d'environ 80 unités de sulphydryle oxydase par kilogramme de farine.
4. Procédé selon la revendication 1, caractérisé en ce que la farine est de la farine de froment.
5. Procédé selon la revendication 1, caractérisé en ce que la sulphydryle oxydase et la glucose oxydase sont présentes dans ladite préparation dans un rapport d'environ 0,35 sur la base des unités de sulphydryle oxydase et de glucose oxydase présentes.

EP 0 321 811 B2

6. Procédé selon la revendication 1, caractérisé en ce que la farine est fortifiée par des conditionneurs de pâte non spécifiques.
7. Procédé selon la revendication 6, caractérisé en ce que la farine est fortifiée avec de l'acide ascorbique en tant que conditionneur de pâte.

5

10

15

20

25

30

35

40

45

50

55